



SKINVIVE™ PRE- + POST-PROCEDURE PATIENT INSTRUCTIONS

PRE-PROCEDURE INSTRUCTIONS

Tell your provider if you are using any medication that can prolong bleeding, such as aspirin, ibuprofen, or other blood thinners. As with any injection, this may increase bruising or bleeding at the injection site.

Tell your provider if you are planning laser treatment, chemical peeling, or any other procedure after SKINVIVE by JUVÉDERM®. There is a possible risk of an inflammatory reaction at the treatment site.

Tell your provider if you are on therapy used to decrease the body's immune response.
Use may result in an increased risk of infection.

Tell your provider if you are pregnant or breastfeeding. The safety for use during pregnancy, or in women who are breastfeeding, has not been studied.

Tell your provider if you have a history of excessive scarring (thick, hard scars). The safety of this product in patients with a history of excessive scarring has not been studied and may result in additional scars.

Tell your provider if you have a history of pigmentation disorders, as use of this product in patients with a history of pigmentation disorders has not been studied and may result in changes in pigmentation.

Stay clear of any blood-thinning medications and supplements for 7 days prior to your appointment to decrease swelling and bruising. Examples include aspirin, ibuprofen, naproxen, vitamin E, and fish oil.

POST-PROCEDURE INSTRUCTIONS

Expect mild swelling, redness, and pin-point bleeding immediately after treatment. You can apply makeup 24 hours after treatment; the injection sites should be virtually undetectable.

Avoid strenuous activity, saunas, gyms, or tanning beds for 24–48 hours post-treatment.

The most commonly reported side effects were redness, lumps/bumps, swelling, bruising, pain, tenderness, firmness, discoloration and itching. Most side effects will resolve within 7 days. If they persist longer, your physician may choose to treat them with medications, such as antibiotics, steroids, or hyaluronidase.

Minimize strenuous exercise, exposure to extensive sun or heat, and alcoholic beverages within the first 24 hours following treatment. Exposure to any of these may cause temporary redness, swelling, and/or itching at the injection site

The SKINVIVE microdroplets purportedly absorb moisture, literally hydrating skin from within. Final results are seen 30-days post injection, and results last up to six months.

If you have questions or concerns after business hours following a procedure, please call us at **802-861-2273** and push **option 5** to be connected to our answering service. **For a true medical emergency, call 911.**

PATIENT SIGNATURE

PRINT NAME

DATE